

Metronidazole Oral Suspension
 Metronidazole 125 mg / 5 ml Oral Suspension

1. NAME OF THE MEDICAL PRODUCT
1.1 Qualitative and Quantitative Composition
 250 ml solution
 Metronidazole 250 mg/ml
 Metronidazole 125 mg/ml

Metronidazole is contained in the form of the active ingredient. The active ingredient is contained in the form of the active ingredient. The active ingredient is contained in the form of the active ingredient. The active ingredient is contained in the form of the active ingredient.

2. CLINICAL PARTICULARS
2.1 Therapeutic Indications

The therapeutic indications of metronidazole are based on its antimicrobial and antiparasitic activity and its mechanism of action. It is used to treat various infections of the oral cavity, including periodontitis, gingivitis, and pharyngitis. It is also used to treat various infections of the gastrointestinal tract, including amoebiasis and trichomoniasis. The drug is also used to treat various infections of the skin and soft tissues, including abscesses and cellulitis.

2.2 Pharmacokinetics and metabolism of metronidazole

Metronidazole is rapidly absorbed from the gastrointestinal tract and is widely distributed in the body. It is excreted in the urine and feces. The half-life of metronidazole is approximately 8 hours. The drug is metabolized in the liver to various active metabolites.

2.3 Contraindications

Metronidazole should not be used in patients with a known hypersensitivity to metronidazole or to any of the excipients.

2.4 Special precautions for use

Patients should be informed of the risks and benefits of metronidazole treatment. Patients should be advised to avoid alcohol during treatment.

2.5 Pregnancy and lactation

Metronidazole should be used with caution in pregnant women and women who are breastfeeding.

2.6 Side effects

The most common side effects of metronidazole are headache, dizziness, and nausea. Other side effects include diarrhea and taste disturbances.

2.7 Overdose

Overdose of metronidazole may cause severe nausea, vomiting, and diarrhea. Treatment is symptomatic.

2.8 Interactions

Metronidazole may interact with other drugs, including alcohol, warfarin, and lithium. Patients should be advised to avoid alcohol during treatment.

2.9 Storage conditions

Metronidazole oral suspension should be stored in a cool, dry place, protected from light.

3. PRESENTATION

Metronidazole oral suspension is available in 250 ml and 500 ml bottles.

4. DATE OF REVISION OF THE TEXT

15/01/2010

2.1 Therapeutic Indications

The therapeutic indications of metronidazole are based on its antimicrobial and antiparasitic activity and its mechanism of action. It is used to treat various infections of the oral cavity, including periodontitis, gingivitis, and pharyngitis. It is also used to treat various infections of the gastrointestinal tract, including amoebiasis and trichomoniasis. The drug is also used to treat various infections of the skin and soft tissues, including abscesses and cellulitis.

2.2 Pharmacokinetics and metabolism of metronidazole

Metronidazole is rapidly absorbed from the gastrointestinal tract and is widely distributed in the body. It is excreted in the urine and feces. The half-life of metronidazole is approximately 8 hours. The drug is metabolized in the liver to various active metabolites.

2.3 Contraindications

Metronidazole should not be used in patients with a known hypersensitivity to metronidazole or to any of the excipients.

2.4 Special precautions for use

Patients should be informed of the risks and benefits of metronidazole treatment. Patients should be advised to avoid alcohol during treatment.

2.5 Pregnancy and lactation

Metronidazole should be used with caution in pregnant women and women who are breastfeeding.

2.6 Side effects

The most common side effects of metronidazole are headache, dizziness, and nausea. Other side effects include diarrhea and taste disturbances.

2.7 Overdose

Overdose of metronidazole may cause severe nausea, vomiting, and diarrhea. Treatment is symptomatic.

2.8 Interactions

Metronidazole may interact with other drugs, including alcohol, warfarin, and lithium. Patients should be advised to avoid alcohol during treatment.

2.9 Storage conditions

Metronidazole oral suspension should be stored in a cool, dry place, protected from light.

3. PRESENTATION

Metronidazole oral suspension is available in 250 ml and 500 ml bottles.

4. DATE OF REVISION OF THE TEXT

15/01/2010

2.1 Therapeutic Indications

The therapeutic indications of metronidazole are based on its antimicrobial and antiparasitic activity and its mechanism of action. It is used to treat various infections of the oral cavity, including periodontitis, gingivitis, and pharyngitis. It is also used to treat various infections of the gastrointestinal tract, including amoebiasis and trichomoniasis. The drug is also used to treat various infections of the skin and soft tissues, including abscesses and cellulitis.

2.2 Pharmacokinetics and metabolism of metronidazole

Metronidazole is rapidly absorbed from the gastrointestinal tract and is widely distributed in the body. It is excreted in the urine and feces. The half-life of metronidazole is approximately 8 hours. The drug is metabolized in the liver to various active metabolites.

2.3 Contraindications

Metronidazole should not be used in patients with a known hypersensitivity to metronidazole or to any of the excipients.

2.4 Special precautions for use

Patients should be informed of the risks and benefits of metronidazole treatment. Patients should be advised to avoid alcohol during treatment.

2.5 Pregnancy and lactation

Metronidazole should be used with caution in pregnant women and women who are breastfeeding.

2.6 Side effects

The most common side effects of metronidazole are headache, dizziness, and nausea. Other side effects include diarrhea and taste disturbances.

2.7 Overdose

Overdose of metronidazole may cause severe nausea, vomiting, and diarrhea. Treatment is symptomatic.

2.8 Interactions

Metronidazole may interact with other drugs, including alcohol, warfarin, and lithium. Patients should be advised to avoid alcohol during treatment.

2.9 Storage conditions

Metronidazole oral suspension should be stored in a cool, dry place, protected from light.

3. PRESENTATION

Metronidazole oral suspension is available in 250 ml and 500 ml bottles.

4. DATE OF REVISION OF THE TEXT

15/01/2010

2.1 Therapeutic Indications

The therapeutic indications of metronidazole are based on its antimicrobial and antiparasitic activity and its mechanism of action. It is used to treat various infections of the oral cavity, including periodontitis, gingivitis, and pharyngitis. It is also used to treat various infections of the gastrointestinal tract, including amoebiasis and trichomoniasis. The drug is also used to treat various infections of the skin and soft tissues, including abscesses and cellulitis.

2.2 Pharmacokinetics and metabolism of metronidazole

Metronidazole is rapidly absorbed from the gastrointestinal tract and is widely distributed in the body. It is excreted in the urine and feces. The half-life of metronidazole is approximately 8 hours. The drug is metabolized in the liver to various active metabolites.

2.3 Contraindications

Metronidazole should not be used in patients with a known hypersensitivity to metronidazole or to any of the excipients.

2.4 Special precautions for use

Patients should be informed of the risks and benefits of metronidazole treatment. Patients should be advised to avoid alcohol during treatment.

2.5 Pregnancy and lactation

Metronidazole should be used with caution in pregnant women and women who are breastfeeding.

2.6 Side effects

The most common side effects of metronidazole are headache, dizziness, and nausea. Other side effects include diarrhea and taste disturbances.

2.7 Overdose

Overdose of metronidazole may cause severe nausea, vomiting, and diarrhea. Treatment is symptomatic.

2.8 Interactions

Metronidazole may interact with other drugs, including alcohol, warfarin, and lithium. Patients should be advised to avoid alcohol during treatment.

2.9 Storage conditions

Metronidazole oral suspension should be stored in a cool, dry place, protected from light.

3. PRESENTATION

Metronidazole oral suspension is available in 250 ml and 500 ml bottles.

4. DATE OF REVISION OF THE TEXT

15/01/2010

